

UNIVERSITY OF COPENHAGEN



## Influence of nutraceuticals on satiety

Astrup, Arne

*Published in:*  
Clinical Nutrition Supplements

*Publication date:*  
2010

*Document version*  
Publisher's PDF, also known as Version of record


*Citation for published version (APA):*  
Astrup, A. (2010). Influence of nutraceuticals on satiety. *Clinical Nutrition Supplements*, 5(1), 2.



**6th Annual Conference of the  
European Nutraceutical Association (ENA)**

March 13th 2010, Vienna, Austria

**Nutraceuticals and Obesity**



**Abstracts**

Publication of this supplement was made possible with support from the European Nutraceutical Association

The European Nutraceutical Association (ENA) is a specialist association devoted to creating a scientific platform for nutraceuticals.  
[www.enaonline.org](http://www.enaonline.org)

disappointing. Many studies employing antioxidants like vitamin E, vitamin C,  $\beta$ -carotin, selenium had no or only marginal effects on the central characteristics of the metabolic syndrome and did not lead to lasting clinical improvements or reduction in the incidence of SX and its follow-up diseases. On the other hand some studies were successful in demonstrating significant improvements following dietary modifications like the mediterranean diet and high intake of fruit and vegetables which are actually the best natural sources of antioxidants. Today it becomes more and more evident that the utilization of complex antioxidant sources like natural fruit and vegetable, red wine, plant oils and even extracts of fruit and vegetable with highly concentrated antioxidants are effective measures to improve the risks of metabolic syndrome and reduce the incidence of consecutive metabolic diseases. Obviously the complexity of antioxidants with many interacting components is the key to the treatment of metabolic oxidative stress diseases like SX.

#### Reference(s)

- [1] Buckland GG, Salas-Salvadó J, Serra-Majem L, Castell C, Cabré J, Salleras-Sanmartí L. Increase in metabolic syndrome as defined by ATPIII from 1992–1993 to 2002–2003 in a Mediterranean population. *Nutr Rev* 2009; 67: S117–25.
- [2] Czernichow S, Couthous A, Bertrais S, Vergnaud AC, Dauchet L, Galan P, Hercberg C. Antioxidant supplementation does not affect fasting plasma glucose in the Supplementation with Antioxidant Vitamins and Minerals (SU.VI.MAX) study in France: association with dietary intake and plasma concentrations. *Am J Clin Nutr* 2006; 84: 395–9.
- [3] Devaraja S, Leonard S, Traber MG, Jialala I. Gamma-tocopherol supplementation alone and in combination with alpha-tocopherol alters biomarkers of oxidative stress and inflammation in subjects with metabolic syndrome. *Free Radic Biol Med* 2008 15; 44(6): 1203–8.
- [4] Ford ES, Mokdad AH, Giles WH, Brown DW. The Metabolic Syndrome and Antioxidant Concentrations Findings From the Third National Health and Nutrition Examination Survey. *Diabetes* 2003; 52: 2346–52.
- [5] Grattagliano G, Palmieri VO, Portincasa P, Moschetta P, Palasciano G. Oxidative stress-induced risk factors associated with the metabolic syndrome: a unifying hypothesis. *J Nutr Biochem* 2008; 19: 491–504.
- [6] Hercberg S, Galan P, Preziosi P, Bertrais S, Mennen L, Malvy D, Roussel A, Favier A, Briançon S. The SU.VI.MAX Study. A randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* 2004; 164: 2335–42.
- [7] Holvoet P, Lee DH, Steffes M, Gross M, Jacobs DR. Association Between Circulating Oxidized Low-Density Lipoprotein and Incidence of the Metabolic Syndrome. *JAMA* 2008; 299(19): 2287–93.
- [8] Nicolson GL. Metabolic syndrome and mitochondrial function: molecular replacement and antioxidant supplements to prevent membrane peroxidation and restore mitochondrial function. *J Cell Biochem* 2007; 100: 1352–69.
- [9] Puchau B, Zulet MA, González de Echávarri A, Hermsdorff HH, Martínez JA. Dietary total antioxidant capacity is negatively associated with some metabolic syndrome features in healthy young adults. *Nutrition* 2009 Sep 25 [Epub ahead of print].

### 3

#### Influence of nutraceuticals on satiety

A. Astrup. *Department of Human Nutrition, Faculty of Life Sciences, University of Copenhagen, Copenhagen, Denmark*

E-mail: ast@life.ku.dk

Obesity has become a global epidemic, and the efforts to curb the rise in prevalence of overweight and obesity have so far not been very successful. The "energy gap" is the daily surplus in energy balance that can explain the average weight gain at a population level, and it has been estimated to be 50–100 kcal. As many dietary factors or substances exert effects on the three components of energy balance, one way to affect the energy gap is to use the inherited properties of these substances. Because the positive energy balance is created by an excessive energy intake (i.e. obese individuals have a higher energy intake than normal weight individuals) many efforts have been exerted to reduce energy intake by reduced portion sizes, reduced energy density foods etc. Recently, more physiological approaches are trying to reduce spontaneous energy intake by enhancement of satiety through naturally occurring nutrients and bioactive food substances working on e.g. gastric emptying, and hormones as ghrelin, CCK, GLP-1, PYY etc.

Clearly, a high intake of dairy protein is reducing spontaneous food intake, and may be one important mechanism to link low-fat dairy and meat products with better weight control. In the large randomised DIogenes dietary intervention study at 8 different European centres, both a higher protein content and lower glycemic index improved retention and weight loss maintenance. Now, research are investigating more specific proteins, peptides and amino acids to obtain a better physiological mechanism of action, and to enable the use of this way to promote satiety without increasing total protein intake.

Dietary calcium reduces fat absorption, and a sufficient intake may also prevent excessive hunger during weight loss diets. Caffeine, chili, mustard, catechins have beneficial effects on energy balance, although the quantitative importance of this may be modest. In conclusion, manipulation of diet composition with an aim to prevent weight gain and weight re-gain is a promising avenue of research.

#### Reference(s)

- [1] A Astrup, M Kristensen, NT Gregersen, A Belza, JK Lorenzen, A Due, TM Larsen. Can bioactive foods impact obesity? The New York Academy of Sciences: "Foods for Health in the 21st Century: A Road Map for the Future." (in press).

### 4

#### Significance of formula diets in obesity therapy

S.C. Bischoff, A. Damms-Machado. *Department of Nutritional Medicine, University of Hohenheim, Stuttgart, Germany*

E-mail: bischoff.stephan@uni-hohenheim.de

The therapeutic spectrum of obesity management comprises first and foremost a conservative approach to achieve weight reduction. Numerous dietary approaches